

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 October 2001 (11.10.2001)

PCT


(10) International Publication Number
WO 01/74812 A1

- (51) International Patent Classification⁷: C07D 417/04, A61K 31/428, 31/5415, A61P 31/04
- (21) International Application Number: PCT/US01/08623
- (22) International Filing Date: 27 March 2001 (27.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/193,631 31 March 2000 (31.03.2000) US
- (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): CISKE, Fred, L. [US/US]; 26431 Riesling Summit, Lawton, MI 49065 (US). GENIN, Michael, J. [US/US]; 33303 Walden Way, Paw Paw, MI 49079 (US).
- (74) Agent: YANG, Lucy, X.; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/74812 A1

(54) Title: NOVEL BENZOSULTAM OXAZOLIDINONE ANTIBACTERIAL AGENTS



(57) Abstract: The present invention provides a compound of formula (1) wherein M is -CH₂- or -CH₂CH₂-, which have potent antibacterial activities.

NOVEL BENZOSULTAM OXAZOLIDINONE ANTIBACTERIAL AGENTS

FIELD OF THE INVENTION

The present invention relates to novel benzosultam oxazolidinones, specifically relates to N-substituted bicyclic benzosultam oxazolidinones; and their preparations. These compounds have potent antibacterial activities.

BACKGROUND OF THE INVENTION

The oxazolidinone antibacterial agents are a novel synthetic class of antimicrobials with potent activity against a number of human and veterinary pathogens, including gram-positive aerobic bacteria such as multiply-resistant staphylococci and streptococci, anaerobic organisms such as bacteroides and clostridia species, and acid-fast organisms such as *Mycobacterium tuberculosis* and *Mycobacterium avium*. The benzosultam oxazolidinones of the present invention may also possess activities against gram-negative organisms such as *Haemophilus influenza* and *Moraxella catarrhalis*.

INFORMATION DISCLOSURE

US Patent No. 5,164,510 discloses 5'-indolinyloxazolidin-2-ones which are useful as antibacterial agents.


US Patent Nos. 5,036,092; 5,036,093; 5,039,690; 5,032,605 and 4,965,268 disclose aminomethyl oxazolidinyl aza cycloalkylbenzene derivatives useful as antibacterial agents.

US Patent Nos. 5,792,765 and 5,684,023 disclose substituted oxazolidinones useful as antibacterial agents.

PCT International Publications WO 98/32438, WO 98/34929, WO 99/36069, WO 9911264, discloses sultam derivatives useful in the treatment of disease states mediated by the chemokine, interleukin-8.

SUMMARY OF THE INVENTION

The present invention provides a compound of formula I



I

- 1 -

or a pharmaceutically acceptable salt thereof wherein:

R^1 is H, NH_2 , NHC_{1-4} alkyl, C_{1-4} alkenyl, OC_{1-4} alkyl, SC_{1-4} alkyl,

$(CH_2)_i-C_{3-6}$ cycloalkyl, or C_{1-4} alkyl, optionally substituted with 1-3 F, 1-2 Cl or CN;

R^2 is H, C_{1-12} alkyl optionally substituted with phenyl or CN, or C_{2-12} alkyl substituted with
5 OH, SH, NH_2 , $-OC_{1-6}$ alkyl, $-NHC_{1-6}$ alkyl, $-NHCOC_{1-6}$ alkyl, $-NHSO_2C_{1-6}$ alkyl,
 $-S(O)_iC_{1-6}$ alkyl, or one to three halo;

Y is O or S;

M is $-(CH_2)_n-$, wherein n is 1 or 2 and

i is 0, 1, or 2.

10 In another aspect, the present invention also provides:

a pharmaceutical composition comprising a compound of formula I or a
pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier,

a method for treating microbial infections in humans or other warm-blooded
animals by administering to the subject in need a therapeutically effective amount of a
15 compound of formula I or a pharmaceutically acceptable salt thereof,

the use of a compound of formula I or a pharmaceutically acceptable salt thereof to
prepare a medicament for treating microbial infections in humans or other warm-blooded
animals, and

The invention also contains novel intermediates and processes that are useful for
20 preparing compounds of formula I.

DETAILED DESCRIPTION OF THE INVENTION

The following definitions are used, unless otherwise described.

The term alkyl, alkenyl, etc. refer to both straight and branched groups, but
25 reference to an individual radical such as "propyl" embraces only the straight chain radical,
a branched chain isomer such as "isopropyl" being specifically referred to.

The carbon atom content of various hydrocarbon-containing moieties is indicated by
a prefix designating the minimum and maximum number of carbon atoms in the moiety,
i.e., the prefix C_{i-j} indicates a moiety of the integer "i" to the integer "j" carbon atoms,
30 inclusive. Thus, for example, C_{1-7} alkyl refers to alkyl of one to seven carbon atoms,
inclusive.

Warm-blooded animals refer to farm animal, companion animal or other type of
animal.

The term "halo" refers to fluoro, chloro, bromo, or iodo

The compounds of the present invention are generally named according to the IUPAC or CAS nomenclature system.

Abbreviations which are well known to one of ordinary skill in the art may be used
5 (e.g. "Ph" for phenyl, "Me" for methyl, "Et" for ethyl, "h" for hour or hours and "rt" for room temperature).

Specific and preferred values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

10 A specific value for R^1 is NH_2 , $-OCH_3$, or C_{1-4} alkyl.

A specific value for R^1 is methyl, ethyl, or isopropyl.

A specific value for R^1 is methyl.

A specific value for R_1 is ethyl.

A specific value for R^2 is C_{1-6} alkyl.

15 A specific value for R^2 is C_{1-6} alkyl substituted with CN.

A specific value for R^2 is benzyl.

A specific value for R^2 is C_{2-6} alkyl substituted with OH, SH, NH_2 , F, $-OC_{1-6}$ alkyl, $-NHC_{1-6}$ alkyl, $-NHCOC_{1-6}$ alkyl, $-NHSO_2 C_{1-6}$ alkyl, $-S(O)_i C_{1-6}$ alkyl, or one to three halo.

A specific value for R^2 is methyl or methyl substituted with CN.

20 A specific value for R^2 is ethyl substituted with fluoro or methoxy.

A specific value for R^2 is $-CH_2CH_2F$.

A specific value for Y is sulfur.

A specific value for Y is oxygen.

A specific value for n is 1.

25 These absolute configurations are called (S)-configuration according to the Cahn-Ingold-Prelog nomenclature system. It will be appreciated by those skilled in the art that compounds of the present invention may have additional chiral centers and be isolated in optically active or racemic form. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic, tautomeric, or stereoisomeric
30 form, or mixture thereof, of a compound of the invention. It is well known in the art how to prepare the optically active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase) and how to

determine activity using the standard tests described herein, or using other similar tests which are well known in the art.


Examples of the present invention are:

- (1) N-([(5S)-3-(1-Methyl-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl)-2-oxo-1,3-oxazolidin-5-yl]methyl)acetamide, (PNU-252307)
- (2) N-([(5S)-3-[1-(2-Fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl]methyl)acetamide, (PNU-254380)
- (3) N-([(5S)-3-[1-(2-Nitriloethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl]methyl)acetamide, (PNU-274919)
- (4) N-([(5S)-3-[1-(2-Methoxyethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl]methyl)acetamide, and (PNU-276461)
- (5) N-([(5S)-3-[1-(2-Fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl]methyl)ethanethioamide. (PNU-254646)

The following describes the preparation of compounds of the present invention. All of the starting materials are prepared by procedures described herein or by procedures that would be well known to one of ordinary skill in organic chemistry.

As shown in CHART I, nitrobenzosultam 1, (can be obtained according to the methods described in J. Het. Chem. 1986, 23, 1645), is first converted to a sodium salt by treatment with a suitable base such as sodium bicarbonate. The nitrogen at the 1-position can then be alkylated by treatment with a variety of alkylating agents including alkyl halides and heating in a suitable solvent such as DMF. These compounds of general structure 2 can be reduced by catalytic hydrogenation in the presence of a suitable catalyst such as palladium on carbon in a suitable solvent such as ethyl acetate, THF, methanol or combinations thereof to afford 5-aminobenzosultams 3. When 3 are treated with magnesium triflate and N-[(2S)oxiranylmethyl] acetamide, prepared by the method of Schaus and Jacobsen (Tetrahedron Lett. 1996, 37, 7937), in a suitable solvent, preferably acetonitrile, the chiral alcohols 4 can be obtained. These compounds can be cyclized to the desired oxazolidinones 5 by reaction with a carbonyl equivalent such as carbonyl diimidazole or preferably N,N'-disuccinimidyl carbonate with an appropriate base such as triethylamine in a mixed solvent system such as acetonitrile/DMF. Additionally, these oxazolidinone amides can be reacted with a sulfurating agent such as Lawesson's Reagent in an appropriate solvent such as THF to obtain the corresponding thioamides 6.


CHART I



In CHART II, the compounds wherein $n = 2$ can be prepared from the known intermediate 7 (Sianesi, E. et al. *Chem. Ber.* 1971, 104, 1880). Nitration of structure 7 provides structure 8. The remaining synthetic steps which lead to structure 9 are similar to the procedures outlined in CHART I.

5

CHART II



It will be apparent to those skilled in the art that the described synthetic procedures are merely representative in nature and that alternative synthetic processes are known to one of ordinary skill in organic chemistry.

The pharmaceutical compositions of this invention may be prepared by combining the compounds of formula I of this invention with a solid or liquid pharmaceutically acceptable carrier and, optionally, with pharmaceutically acceptable adjuvants and excipients employing standard and conventional techniques. Solid form compositions include powders, tablets, dispersible granules, capsules, cachets and suppositories. A solid carrier can be at least one substance which may also function as a diluent, flavoring agent, solubilizer, lubricant, suspending agent, binder, tablet disintegrating agent, and encapsulating agent. Inert solid carriers include magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, cellulosic materials, low melting wax, cocoa butter, and the like. Liquid form compositions include solutions, suspensions and emulsions. For example, there may be provided solutions of the compounds of this invention dissolved in water and water-propylene glycol and water-polyethylene glycol systems, optionally containing suitable conventional coloring agents, flavoring agents, stabilizers and thickening agents.

Preferably, the pharmaceutical composition is provided employing conventional techniques in unit dosage form containing effective or appropriate amounts of the active component, that is, the compounds of formula I according to this invention.

The quantity of active component, that is the compound of formula I according to this invention, in the pharmaceutical composition and unit dosage form thereof may be varied or adjusted widely depending upon the particular application, the potency of the


magnesium trifluoromethanesulfonate (0.42g, 1.3mmol) in dry CH_3CN (10mL) at room temperature. After 20hrs solvent is evaporated and the residue purified by chromatography (4%MeOH/ CH_2Cl_2) to give a foamy solid (0.23g, 79%). HRMS (FAB) calcd for $\text{C}_{13}\text{H}_{19}\text{N}_3\text{O}_4\text{S} + \text{H}_1$ 314.1174, found 314.1174.

- 5 Step: 3 Preparation of N-([(5S)-3-(1-methyl-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl)-2-oxo-1,3-oxazolidin-5-yl)methyl]acetamide PNU-252307

The previous product (0.20g, 0.63mmol) is dissolved in CH_3CN (5mL) and DMF (2.5mL). *N,N'*-Disuccinimidyl carbonate (0.23g, 0.90mmol) is added followed by triethylamine (0.26mL, 1.9mmol) and the mixture stirred at room temperature for 20hrs.

- 10 The mixture is poured into CH_2Cl_2 (30mL) and washed with H_2O (3x20mL). The organics are dried (Na_2SO_4), filtered and solvent evaporated. The residue is chromatographed (3%MeOH/ CH_2Cl_2) to obtain a white solid (0.142g, 66%). Mp 84-6 °C. HRMS (FAB) calcd for $\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}_5\text{S} + \text{H}_1$ 340.0967, found 340.0965.

- 15 Example 2 Preparation of N-([(5S)-3-[1-(2-fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl]acetamide PNU-254380



- Step 1: Preparation of 1-(2-fluoroethyl)-5-nitro-1,3-dihydro-2H-2,1-benzisothiazole-2,2-dione

NaHCO_3 (0.93g, 11.0mmol) is dissolved in H_2O (15mL) and 5-Nitro-1,3-dihydro-2H-2,1-benzisothiazole-2,2-dione (1) (1.90g, 8.9mmol) is added with stirring. The mixture is heated to 80°C for 0.5 hrs and a yellow solid formed. The mixture is cooled to 0°C and filtered. The solid is washed with cold H_2O (15mL) then with cold EtOH (25mL). The

- 25 yellow solid thus obtained is dried under high vacuum then dissolved in dry DMF (15mL).

1-Bromo-2-fluoroethane (1.52 mL, 20.0mmol) is added and the solution heated to 100°C for 4 hrs. The solution is cooled to room temperature then poured into ice water (50mL).

The solid is collected by filtration and gave crystals (1.55g, 67%) after recrystallization

from EtOH. Mp 142-4 °C. HRMS (FAB) calcd for $C_9H_9FN_2O_4S + Na_1$ 283.0165, found 283.0166.


Step 2: Preparation of 5-amino-1-(2-fluoroethyl)-1,3-dihydro-2H-2,1-benzisothiazole-2,2-dione

5 The product of Step 1 (1.25g, 4.8mmol) is dissolved into EtOAc (30mL) in a Parr bottle and 10% Pd/C (100mg) added under nitrogen. The mixture is hydrogenated on a Parr apparatus for 4 hrs at 40psi. Filtration and evaporation of solvent gave a solid (1.1g, 99%). Mp 119-21 °C. HRMS (FAB) calcd for $C_9H_{11}FN_2O_2S + H_1$ 231.0603, found 231.0610.

Step 3: N-((5S)-3-[1-(2-fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-10 5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide PNU-254380

The previous 5-amino product (0.95g, 4.1mmol) is added to a mixture of N-[(2S)oxiranylmethyl] acetamide (0.81g, 7.0mmol) and magnesium trifluoromethanesulfonate (2.0g, 6.2 mmol) in dry CH_3CN (45mL) at room temperature. After 20hrs solvent is evaporated and the residue chromatographed (4% MeOH/ CH_2Cl_2).
15 The intermediate product is dissolved in CH_3CN (25mL) and DMF (12mL). N,N'-Disuccinimidyl carbonate (2.25g, 9.0mmol) is added followed by triethylamine (2.6mL, 18.8mmol) and the mixture stirred at room temperature for 20hrs. The mixture is poured into CH_2Cl_2 (30mL) and washed with H_2O (3x20mL). The organics are dried (Na_2SO_4), filtered and solvent evaporated. The residue is chromatographed (3%MeOH/ CH_2Cl_2) to
20 obtain a white solid (0.58g, 38%). Mp 84-7 °C (dec). HRMS (FAB) calcd for $C_{15}H_{18}FN_3O_5S + H_1$ 372.1029, found 372.1021.

Example 3 Preparation of N-((5S)-3-[1-(2-nitriloethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide PNU-274919



25

Step 1: Preparation of 2-(5-nitro-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-1-yl)acetonitrile

$NaHCO_3$ (1.48g, 17.6mmol) is dissolved in H_2O (20mL) and 5-Nitro-1,3-dihydro-2H-2,1-benzisothiazole-2,2-dione (1) (3.17g, 14.8mmol) is added with stirring. The mixture is


heated to 80°C for 0.5 hrs and a yellow solid formed. The mixture is cooled to 0°C and filtered. The solid is washed with cold H₂O (25mL) then with cold EtOH (45mL). The yellow solid thus obtained (intermediate sodium salt) is dried under high vacuum. A portion of this material (1.0g, 4.2mmol) is dissolved in dry DMF (7.0mL).

- 5 Bromoacetonitrile (0.34mL, 5.0mmol) is added and the solution heated to 100°C for 2 hrs. The solution is cooled to room temperature then poured into ice water (50mL). The solid is collected by filtration and gave an off-white solid (0.73g, 69%) after recrystallization from EtOH. Mp 200-2 °C. HRMS (FAB) calcd for C₉H₇N₃O₄S +H₁ 254.0235, found 254.0235.

Step 2: N-((5S)-3-[1-(2-nitriloethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide PNU-274919

- 10 The product of Step 1 (0.73g, 2.9mmol) is dissolved into EtOAc (30mL) in a Parr bottle and 10% Pd/C (100mg) added under nitrogen. The mixture is hydrogenated on a Parr apparatus for 2 hrs at 40psi. Filtration and evaporation of solvent gave a yellow solid which is added to a mixture of N-[(2S)oxiranylmethyl]acetamide (0.97g, 8.0mmol) and
- 15 magnesium trifluoromethanesulfonate (1.3g, 4.0 mmol) in dry CH₃CN (30mL) at room temperature. After 20hrs solvent is evaporated and the residue chromatographed (4% MeOH/CH₂Cl₂). The intermediate product is dissolved in CH₃CN (20mL) and DMF (10mL). N,N'-Disuccinimidyl carbonate (1.0g, 4.1mmol) is added followed by triethylamine (1.2mL, 8.7mmol) and the mixture stirred at room temperature for 20hrs. The
- 20 mixture is poured into CH₂Cl₂ (30mL) and washed with H₂O (3x20mL). The organics are dried (Na₂SO₄), filtered and solvent evaporated. The residue is chromatographed (3%MeOH/CH₂Cl₂) to obtain a pale yellow solid (0.12g, 11%). Mp 124-6 °C (dec). HRMS (FAB) calcd for C₁₅H₁₆N₄O₅S +H₁ 365.0919, found 365.0915.

- 25 **Example 4** Preparation of N-((5S)-3-[1-(2-methoxyethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide PNU-276461




Step 1: 1-(2-methoxyethyl)-5-nitro-1,3-dihydro-2H-2,1-benzisothiazole-2,2-dione

NaHCO₃ (1.48g, 17.6mmol) is dissolved in H₂O (20mL) and 5-Nitro-1,3-dihydro-2H-2,1-benzisothiazole-2,2-dione (1) (3.17g, 14.8mmol) is added with stirring. The mixture is heated to 80°C for 0.5 hrs and a yellow solid formed. The mixture is cooled to 0°C and
5 filtered. The solid is washed with cold H₂O (25mL) then with cold EtOH (45mL). The yellow solid thus obtained (intermediate sodium salt) is dried under high vacuum. A portion of this material (1.0g, 4.2mmol) is dissolved in dry DMF (7.0mL). 2-Bromoethyl methyl ether (1.41mL, 15.0mmol) and potassium iodide (10mg) are added and the solution heated to 130°C for 5 days. The solution is cooled to room temperature then poured into
10 ice water (50mL). The solid is collected by filtration and gave a solid (0.77g, 67%) after chromatography (50% EtOAc/ Heptane). Mp 130-2 °C. HRMS (FAB) calcd for C₁₀H₁₂N₂O₅S +H₁ 273.0545, found 273.0548.

Step 2: N-((5S)-3-[1-(2-methoxyethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide PNU-276461

15 The product of Step 1 (0.70g, 2.6mmol) is dissolved into EtOAc (30mL) in a Parr bottle and 10% Pd/C (100mg) added under nitrogen. The mixture is hydrogenated on a Parr apparatus for 2 hrs at 40psi. Filtration and evaporation of solvent gave a residue which is added to a mixture of N-[(2S)oxiranylmethyl]acetamide (0.115g, 1.0mmol) and magnesium trifluoromethanesulfonate (0.32g, 1.0 mmol) in dry CH₃CN (10mL) at room temperature.
20 After 20hrs solvent is evaporated and the residue is dissolved in CH₃CN (10mL) and DMF (5mL). N,N'-Disuccinimidyl carbonate (0.30g, 1.2mmol) is added followed by triethylamine (0.34mL, 2.4mmol) and the mixture stirred at room temperature for 20hrs. The mixture is poured into CH₂Cl₂ (30mL) and washed with H₂O (3x20mL). The organics are dried (Na₂SO₄), filtered and solvent evaporated. The residue is chromatographed
25 (3%MeOH/CH₂Cl₂) to obtain a pale yellow foamy solid (0.114g, 38%). HRMS (EI) calcd for C₁₆H₂₁N₃O₆S 383.1151, found 383.1149.


Example 5 Preparation of N-((5S)-3-[1-(2-fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)ethanethioamide PNU-254646



The product from example 2 (0.195g, 0.52mmol) and Lawesson's Reagent (0.21g, 0.52mmol) are stirred in dry THF (20mL) and heated to reflux for 20 hrs. The mixture is cooled to room temperature and solvent evaporated. The residue is chromatographed
5 (4%MeOH/CH₂Cl₂) to give a solid (0.178g, 88%). Mp 90-3 °C (dec). HRMS (FAB) calcd for C₁₅H₁₈FN₃O₄S₂ +H₁ 388.0801, found 388.0805.

CLAIMS

1. A compound of formula I



- 5 I
or a pharmaceutically acceptable salt thereof wherein:

R¹ is

- a) H,
b) NH₂,
10 c) NHC₁₋₆ alkyl,
d) C₁₋₆ alkenyl,
e) OC₁₋₆ alkyl, SC₁₋₆ alkyl,
f) (CH₂)_i-C₃₋₆ cycloalkyl, or
g) C₁₋₆ alkyl, optionally substituted with one to three halo;
15 R² is
a) H,
b) C₁₋₁₂ alkyl, optionally substituted with phenyl or CN, or
c) C₂₋₁₂ alkyl substituted with OH, SH, NH₂, -OC₁₋₆ alkyl, -NHC₁₋₆ alkyl,
-NHCOC₁₋₆ alkyl, -NHSO₂C₁₋₆ alkyl, -S(O)_iC₁₋₆ alkyl, or one to three halo;
20 Y is O or S;
M is -(CH₂)_n-, wherein n is 1 or 2; and
i is 0, 1, or 2.

2. A compound of claim I wherein R¹ is NH₂, -OCH₃, or C₁₋₄ alkyl.
25
3. A compound of claim I wherein R¹ is methyl
4. A compound of claim I wherein R¹ is ethyl.
5. A compound of claim I wherein R² is C₁₋₆ alkyl, optionally substituted with phenyl
30 or CN.

6. A compound of claim I wherein R² is C₂₋₆ alkyl substituted with OH, SH, NH₂, F, -OC₁₋₆ alkyl, -NHC₁₋₆ alkyl, -NHCOC₁₋₆ alkyl, -NHSO₂ C₁₋₆ alkyl, or -S(O)_iC₁₋₆ alkyl.
- 5
7. A compound of claim I wherein R² is C₁₋₄ alkyl optionally substituted with CN, or C₂₋₄ alkyl substituted with fluoro or OC₁₋₄ alkyl.
8. A compound of claim I wherein R¹ is C₁₋₄ alkyl; R² is C₁₋₄ alkyl optionally substituted with CN, or C₂₋₄ alkyl substituted with fluoro or OC₁₋₄ alkyl; Y is sulfur or oxygen; and n is 1.
- 10
9. A compound of claim 8 wherein Y is sulfur.
- 15
10. A compound of claim 8 wherein R¹ is CH₃ or CH₃CN; R² is CH₂CH₂F or CH₂CH₂OCH₃; Y is sulfur or oxygen, n is 1.
11. A compound of claim 10 wherein Y is sulfur.
- 20
12. A compound of claim 1 which is
- (a) N-(((5S)-3-(1-Methyl-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl)-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide,
- (b) N-(((5S)-3-[1-(2-Fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide,
- 25 (c) N-(((5S)-3-[1-(2-Nitriloethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide,
- (d) N-(((5S)-3-[1-(2-Methoxyethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)acetamide, or
- (e) N-(((5S)-3-[1-(2-Fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)ethanethioamide.
- 30
13. A compound of claim 1 which is N-(((5S)-3-[1-(2-Fluoroethyl)-2,2-dioxo-2,3-dihydro-1H-2,1-benzisothiazol-5-yl]-2-oxo-1,3-oxazolidin-5-yl)methyl)ethanethioamide.

14. A use of compounds of claims 1 through 13 for manufacturing of medicinals for treating microbial infections.

15. The use of claim 14 wherein said medicinal is administered orally, parenterally, transdermally, or topically in a pharmaceutical composition.

16. The use of claim 14 wherein said medicinal is administered in an amount of from about 0.1 to about 500 mg/kg of body weight/day.

10 17. The use of claim 14 wherein said medicinal is administered in an amount of from about 1 to about 50 mg/kg of body weight/day.

18. A use for treating microbial infections of claim 14 wherein the infection is skin infection.

15

19. A use for treating microbial infections of claim 14 wherein the infection is eye infection.

20. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

20

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/08623

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/04 A61K31/428 A61K31/5415 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 164 510 A (BRICKNER S J) 17 November 1992 (1992-11-17) cited in the application the whole document, particularly example 148	1-20
Y	EP 0 738 726 A (BAYER AG) 23 October 1996 (1996-10-23) the whole document, particularly example 102	1-20
Y	WO 99 37641 A (BAYER AKTIENGESELLSCHAFT) 29 July 1999 (1999-07-29) the whole document, particularly example 107	1-20



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document delining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

9 July 2001

Date of mailing of the international search report

19/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Allard, M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/08623

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5164510	A	17-11-1992	US 5182403 A	26-01-1993
			US 5225565 A	06-07-1993
			AT 112773 T	15-10-1994
			AT 201870 T	15-06-2001
			AU 617871 B	05-12-1991
			AU 4195789 A	02-04-1990
			CA 1335103 A	04-04-1995
			DE 68918792 D	17-11-1994
			DK 45591 A	13-03-1991
			EP 0359418 A	21-03-1990
			EP 0434714 A	03-07-1991
			EP 0609905 A	10-08-1994
			HK 1002234 A	07-08-1998
			JP 11080139 A	26-03-1999
			JP 2865211 B	08-03-1999
			JP 4500665 T	06-02-1992
			KR 138262 B	15-05-1998
			WO 9002744 A	22-03-1990
EP 738726	A	23-10-1996	DE 19544106 A	24-10-1996
			AU 705071 B	13-05-1999
			AU 5073596 A	31-10-1996
			BG 100525 A	31-03-1997
			BR 9602016 A	07-04-1998
			CA 2174473 A	22-10-1996
			CN 1138582 A	25-12-1996
			CZ 9601142 A	16-07-1997
			HR 960159 A	31-08-1997
			HU 9601001 A	28-04-1998
			JP 8301869 A	19-11-1996
			NO 961559 A	22-10-1996
			NZ 286400 A	27-04-1998
			SG 52774 A	28-09-1998
			SK 50096 A	07-05-1997
			TR 960963 A	21-11-1996
			US 6069160 A	30-05-2000
			ZA 9603138 A	04-11-1996
WO 9937641	A	29-07-1999	DE 19802239 A	29-07-1999
			AU 2616199 A	09-08-1999
			EP 1049692 A	08-11-2000